

LA-UR-18-27741

Approved for public release; distribution is unlimited.

Title: Investigations in Cell-Centered Hydrodynamics

Author(s): Green, Lauren Maranne
Thorneloe, Logan Alexander

Intended for: XCP Workshop Quick Look

Issued: 2018-08-14

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



Investigations in Cell-Centered Hydrodynamics

Computational Physics Workshop



Lauren Green and Logan Thorneloe


Mentor: Marc Charest



Operated by Los Alamos National Security, LLC for the U.S. Department of Energy's NNSA

Logan Thorneloe

- **Attending Brigham Young University, Provo**
- **Bachelor of Science, Computer Engineering**
- **Personal Interests: water polo, swimming, technology**
- **Research Interests: magnetic resonance physics, bioelectronics, machine learning, computers**



Lauren Green

- BS in Nuclear Engineering and Radiological Sciences (NERS), Class of 2018
- starting NERS MSE this fall
- personal interests: travel, food, puppy




Comparison of CCH Lagrangian Schemes

Nodal velocity computation via sub-cell force balance:

$$U_p^n = (M_p^n)^{-1} \sum_{c \in C(p)} [l_{pc}^n P_c(\mathbf{x}_p^n) \mathbf{n}_{pc}^n + M_{pc}^n U_c(\mathbf{x}_p^n)]$$

$$M_p^n = \sum_{c \in C(p)} M_{pc}^n$$

Corner matrices corresponding to GLACE[2], EUCCLHYD[2], and Burton et al.[1] schemes




$$M_{pc}^{GLACE} = z_c l_{pc} (\mathbf{n}_{pc} \otimes \mathbf{n}_{pc})$$

$$M_{pc}^{EUCCL} = z_c [l_{pc}^- (\mathbf{n}_{pc}^- \otimes \mathbf{n}_{pc}^-) + l_{pc}^+ (\mathbf{n}_{pc}^+ \otimes \mathbf{n}_{pc}^+)]$$

$$M_{pc}^{BURT^1} = z_c l_{pc}^- |\mathbf{n}_{pc}^- \cdot \hat{\mathbf{u}}_c| + z_c l_{pc}^+ |\mathbf{n}_{pc}^+ \cdot \hat{\mathbf{u}}_c|$$

Numerical Tests in 2D Cartesian Geometry

Checkerboard Problem:





GLACE

EUCCLHYD


Conclusions

- GLACE scheme experiences instabilities in both checkerboard and sedov problem
- EUCCLHYD performs well in both test cases, most accurate of the three schemes
- Burton et al.'s model is more dissipative and improves mesh robustness, but smaller time step


Sedov Problem:



GLACE



EUCCLHYD




BURT. et al.

Linear Least Squares Reconstruction

Goal: better estimate nodal velocity


- utilizes values at centroids to construct gradient field for a single cell



- use gradient to calculate values at cell vertices

$$u = u_0 + \frac{du}{dx}(x - x_0) + \frac{du}{dy}(y - y_0)$$

As the number of cells in a mesh increases, the error of least squares function decreases



References

- [1] D.E. Burton, T.C. Carney, N.R. Morgan, S.K. Sambasivan, M.J. Shashkov. A cell-centered Lagrangian Godunov-like method for solid-dynamics. *Computers and Fluids*, 2012.
- [2] P.-H. Maire. Contribution to the numerical modeling of Inertial Confinement Fusion, 2011. Chapter 3, pp 57-158.
- [3] Timothy J. Barton. Recent Developments in High Order K-Exact Reconstruction of Unstructured Meshes. *31st Aerospace Sciences Meeting & Exhibit*, 1993.